arXiv:1303.0530v2 [astro-ph.HE] 4 Apr 2013

AMS-02 positron excess: new bounds on dark matter models and hint for primary electron spectrum hardening

Lei Feng,[1] Rui-Zhi Yang,[1,2] Hao-Ning He,[1] Tie-Kuang Dong,[1] Yi-Zhong Fan*,[1] and Jin Chang[1]
[1]*Key Laboratory of Dark Matter and Space Astronomy,*
Purple Mountain Observatory, Chinese Academy of Sciences, Nanjing 210008, China
[2]*University of Chinese Academy of Sciences, Beijing, 100012, China*

The bounds on dark matter models have been updated with the first result of AMS-02 on positron fraction in cosmic rays. The channels of annihilation/decay into $\tau^+\tau^-$ pairs have been excluded by the Fermi-LAT Galactic diffuse emission data. The channel $\chi\chi \to \mu^+\mu^-$ is allowed only for very cored dark matter distribution model (e.g., isothermal-sphere profile) while $\chi \to \mu^+\mu^-$ is viable. Interestingly, at energies ≤ 100 GeV a positron/electron component from either dark matter annihilation/decay or pulsar seems to be favored by the AMS-02 data. In the case of annihilation of dark matter with a rest mass ~ 100 GeV, the cross section into electron/positron pairs is $\sim 10^{-26}$ cm^3 s^{-1}. We also suggest that the primary electron spectrum may display hardening at ~ 240 GeV, as observed by ATIC, CREAM and PAMELA in the cosmic-ray-nuclei spectra. Depending on the degree of the spectrum hardening, the subsequent positron-to-electron ratio can get flattened or even decreased, which may account for the flattening of positron fraction at energies $\gtrsim 200$ GeV discovered by AMS-02.

PACS numbers: 95.35.+d, 96.50.S-, 97.60.Jd, 98.38.Mz

I. INTRODUCTION

In the standard cosmic ray model, most cosmic ray electrons are from supernova remnants while the cosmic ray positrons are mainly produced through hadronic processes as cosmic ray protons collide with intergalactic hydrogen [1, 2]. In conventional approach, the injection spectrum of electrons (positrons) is taken as a single power-law. Since diffusion and electron/positron cooling are more efficient in higher energies, the spectrum should soften with energy and the positron-to-electron ratio (i.e., $\Phi_{e^+}/(\Phi_{e^+} + \Phi_{e^-})$, where Φ is the flux) should drop with energy monotonously [3]. Hence there should be no prominent feature at TeV energies in cosmic ray electron/positron total spectrum, neither in the positron-to-electron ratio. The situation has changed dramatically since 2008. The Advanced Thin Ionization Calorimeter (ATIC [4]), Polar Patrol Balloon-borne Electron Telescope with Scintillating Fibers (PPB-BETS [5]), The High Energy Stereoscopic System (HESS [6]) and Fermi Large area Telescope (Fermi-LAT [7]) reported cosmic ray electron/positron total spectrum up to TeV and found a hardening/bump in the energy range from 100 GeV to 1 TeV. Almost simultaneously the Payload for Antimatter Matter Exploration and Light-nuclei Astrophysics (PAMELA) discovered an unambitious rise of the positron-to-electron ratio above ~ 10 GeV [8]. Such a peculiar rising behavior has been confirmed by Fermi-LAT though in an indirect way [9]. People call the well-established electron/positron total spectrum hardening/bump and the increasing positron-to-electron ratio, unexpected in the standard model, the excesses or anomalies. These interesting features draw a lot of attention, and various physical origins, in particular new astrophysical sources and new physics (dark matter), have been extensively explored [see 10, for recent reviews]. Very recently, the AMS-02 collaboration has released their first result on positron fraction in cosmic rays, which confirms the positron excess with unprecedented accuracy up to the energy ~ 350 GeV [11].

On the other hand, the protons and Helium are the most abundant cosmic-ray components, and the spectrum of these cosmic-rays up to the so-called “knee” can be described by a single power law [1, 2]. Surprisingly, the spectra of protons (Helium) measured by ATIC [12], Cosmic Ray Energetics And Mass (CREAM [13]) and PAMELA [14] show a remarkable hardening at the magnetic rigidity $\sim$ several hundred GV (GeV/nucleon). We call such a kind of spectrum hardening the cosmic ray nuclei excesses, which challenge the current paradigm of cosmic-ray acceleration and propagation in the Galaxy. Various interpretations have been put forward, such as the spectrum superposition of the local source and the background [15], the non-linear acceleration of cosmic rays [16], and superposition of the injection spectra of the cosmic ray sources [17]. Instead of performing an advanced study of the possible physical origin of the nuclei excesses, in this work we simply take such observational indication (i.e., the hardening of the cosmic ray proton and helium spectra above ~ 240 GV [14]) as the main motivation to consider the possibility that the primary electron spectrum gets hardened at ~ 240 GeV.

This work is structured as follows. In section II we take the latest AMS-02 positron ratio data to set new bounds on the dark matter models. In section III we discuss the possible positron-to-electron ratio signal in the presence of primary electron spectrum hardening and then compare with the observational data. The implication on the physical origin of the cosmic ray nuclei excesses is also investigated. We summarize our results with some discussion in Sec. IV.

II. AMS-02 POSITRON RATIO DATA: BOUNDS ON DARK MATTER MODELS

Dark matter is a form of matter necessary to account for gravitational effects observed in very large scale structures such as the flat rotation curves of galaxies and the gravitational lensing of light by galaxy clusters that cannot be accounted for by the amount of observed/normal matter [18]. The most widely discussed candidate is the so-called weakly interacting massive particles (WIMPs), which may annihilate with each other or decay and then produce particle pairs such as photons, electrons and positrons and so on [18]. That is why dark matter may be able to account for the observed positron and electron excesses, as extensively examined in the literature [19–21].

The high quality AMS-02 data confirms the positron excess discovered by PAMELA with unprecedented accuracy in the energy range of $10-350$ GeV [11]. It is the time to explore the allowed WIMP parameter regions in both the annihilation model and the decay model. The electrons originated from dark matter annihilation/decay will suffer from inverse Compton scattering of interstellar background photons (e.g., cosmic microwave background, dust emission and star light) and boost these photons to GeV energies, becoming part of the Galactic diffuse emission. Hence the Galactic diffuse emission detected by space telescopes can be used to reliably constrain the physical parameters of dark matter particles [18]. The latest bounds set by the Fermi-LAT Galactic diffuse emission data are presented in [22]. To make use of such bounds, our dark matter density distribution profiles as well as the cosmic ray diffusion parameters are taken to be the same as that adopted in [22]. The smooth dark matter density ρ are parameterized with a Navarro-Frenk-White (NFW) spatial profile [23]

$$\rho(r) = \rho_0 (1 + r_\odot/r_{\rm s})^2/[(1 + r/r_{\rm s})^2 r/r_\odot], \tag{1}$$

and the isothermal-sphere (ISO) profile [24]

$$\rho(r) = \rho_0 (r_\odot^2 + r_{\rm c}^2)/(r^2 + r_{\rm c}^2), \tag{2}$$

respectively. Where $\rho_0 = 0.43$ GeV cm^{-3} is the local density of dark matter, $r_\odot = 8.5$ kpc is the distance of the solar system from the Galactic center, $r_{\rm s} = 20$ kpc and $r_{\rm c} = 2.8$ kpc are two scale radii.



FIG. 1: The regions of parameter space (blue and green regions, 99.5% and 99.999% confidence levels) which provide a reasonable fit to the positron fraction data of AMS-02 [11] comparing with the bounds set by Fermi-LAT Galactic diffuse emission (adopted from [22], the black lines represent the upper limits on $\langle\sigma v\rangle$ found in the analysis with no model of the astrophysical background, while the red lines are the bounds found in the analysis with a modeling of the background). In each panel, the annihilation/decay channel as well as the dark matter distribution model have been marked. If one takes the bounds found in the analysis with a modeling of the background of [22] (i.e., the red lines), the channel $\chi\chi \to \mu^+\mu^-$ is allowed only for the isothermal-sphere like dark matter distribution model and the decay channel $\chi \to \mu^+\mu^-$ is viable. We do not present the cases of annihilation/decay into $\tau^+\tau^-$ since they have been excluded (by the red lines).

In the modeling, we consider the channels of annihilation/decay into $\mu^+\mu^-$ and $\tau^+\tau^-$. These particles propagate in the Galaxy. In general, it is difficult to solve the propagation equation analytically, given the complicated distributions of the sources, interstellar matter, radiation field and magnetic field. In this work we adopt the GALPROP [25] package to numerically calculate the propagation of the cosmic ray particles, including the contribution from dark matter annihilation or decay. The cosmic ray diffusion parameters are the following, including the Halo height $z_{\rm h} = 4$ kpc, the diffusion coefficient $D_0 = 5.3 \times 10^{28}$ cm^2 s^{-1}, the diffusion index $\delta = 1/3$, the Alfven velocity $V_{\rm A} = 33.5$ km s^{-1}, the nucleon injection indexes below and above the break rigidity $\rho_{\rm br} = 11.5$ GV are 1.88 and 2.39, respectively.

In Fig.1 we present the parameter regions allowed by the latest AMS-02 data, where $\langle\sigma v\rangle$ ($\tau_{\rm dec}$) is the velocity-weighted-cross-section (lifetime) of dark matter annihilation (decay) into $\mu^+\mu^-$. Our parameter space regions are much more compact than that obtained in [21] since the AMS-02 data have much smaller errors and extend to much higher energies. Evidently, if one takes the bounds found in the analysis with a modeling of the background of [22] (i.e., the red lines), the channel $\chi\chi \to \mu^+\mu^-$ is allowed only for the isothermal-sphere like dark matter distribution model but the decay channel $\chi \to \mu^+\mu^-$ is still viable. The channels of annihilation/decay into $\tau^+\tau^-$ are not presented since they have been excluded by the Galactic diffuse emission data.

Alternatively, the positron excess detected by PAMELA/Fermi-LAT/AMS-02 may be mainly contributed by pulsars. It is widely assumed that the high energy electrons/positrons are generated through the cascade of electrons accelerated in the magnetosphere of pulsars [28, 29]. The energy spectrum of e^+e^- injected to the galaxy from pulsars can be parameterized as a broken power-law with the cutoff at E_c, i.e., ${\rm d}N/{\rm d}E \propto A_{\rm psr} E^{-\alpha} \exp(-E/E_c)$, where E_c ranges from several tens GeV to higher than TeV and the power-law



FIG. 2: The blue (red) solid line represents the best fit to the positron ratio data in the "background + pulsar" ("background + dark matter + pulsar") model. For the three-component model, we also show the positron-to-electron ratio (at energies > 240 GeV) expected in different primary electron spectrum models: the solid line is for a smooth spectrum without hardening, while the dashed and dotted lines are for the spectra with hardening at the energy $E = 240$ GeV (the indexes of hardened primary electron spectra are marked in the Figure). The AMS-02 data are taken from [11].

index α ranges from 1 to 2.2 depending on the gamma-ray and radio observations [28–30]. Following [32], $E_c = 860$ GeV is adopted in the modeling. The spatial distribution of pulsars is parameterized as [31]

$$f(R,z) \propto \left(\frac{R}{R_\odot}\right)^a \exp\left[-\frac{b(R-R_\odot)}{R_\odot}\right] \exp\left(-\frac{|z|}{z_s}\right), \tag{3}$$

where $R_\odot = 8.5$ kpc is the distance of solar system from the Galactic center, $z_s \approx 0.2$ kpc is the scale height of the pulsar distribution, $a = 2.35$ and $b = 5.56$. The other parameter appearing in the above equation is the normalization factor $A_{\rm psr}$, and it will be determined in our modeling. Following [27], we consider the two-component (background + pulsar) model as well as the three-component (background + dark matter + pulsar) model. The background electron injection indexes below and above the break rigidity $\rho_{\rm br,e} = 4$ GV are taken to be 1.54 and 2.6, respectively. For the two-component (three-component) model our best fit of the data at energies ≥ 10 GeV gives a minimum $\chi^2 = 53.35$ (27.59), where the parameter $\alpha = 1.28$ (1.265) is adopted. Please see the solid lines in Fig.2 for the best fits. Such a result seems to favor the three-component model, in which the annihilation cross section $<\sigma v>_{\chi\chi\rightarrow e^+e^-} \sim 10^{-26}$ cm^3 s^{-1} as well as the dark matter rest mass $m_\chi \sim 100$ GeV are needed. As noted in [27], such a cross section is consistent with all of the current bounds of the indirect detection measurements.

III. POSITRON-TO-ELECTRON RATIO BEHAVIOR: THE CASE OF PRIMARY-ELECTRON SPECTRUM HARDENING AT ~ 240 GEV

ATIC, CREAM and PAMELA all found considerable cosmic-ray-nuclei spectrum hardening above the magnetic rigidity ~ 240 GV. One would naturally speculate that the primary electron spectrum also gets hardened at ~ 240 GeV. With the given electron/positron total spectrum detected by (for example) Fermi-LAT, the hardening of primary electron component inevitably leads to the softening of the rest "additional component" which has been widely assumed to consist of electron and positron pairs. Hence the positron-to-electron ratio should be different from that in the absence of a spectrum hardening. Such a possible modification is of our interest. The change of the spectral index of primary electrons at ~ 240 GeV is hard to reliably estimate and we assume it is within the range $0.18-0.3$, consistent with the proton (helium) spectrum hardening found by PAMELA [14]. The following calculation is straightforward. Firstly, for illustration, we take the analytical approach. As shown in the Fig.23 of [26], the primary electron spectrum at 240 GeV is $\propto E^{-3.35}$ (see also Fig.1 of [27] for a rather similar result) and the total electron/positron spectrum is $\propto E^{-3.2}$ for $E > 240$ GeV. Therefore, if the primary electron spectrum gets hardened by $\sim E^{0.18}$ ($E^{0.3}$) at $E \geq 240$ GeV, the subsequent positron-to-electron ratio would get flattened (decreased)[1]. Secondly, to check our speculation more qualitatively, we perform a joint fit to the AMS-02 positron-to-electron data [33] and the Fermi-LAT electron/positron total spectrum data [26] and then evaluate the positron-to-electron ratio beyond the PAMELA energy range in the cases with and without primary electron spectrum hardening. The numerical results are presented in Fig.2. As expected in our analytical approach, the resulting positron-to-electron ratio at energies above ~ 240 GeV gets flattened or decreased, depending on the degree of the primary electron spectrum hardening. The current AMS-02 data show tentative evidence for flattening at energies above ~ 260 GeV. However the errors are still large and a robust conclusion is not possible. A full test of our prediction by AMS-02 may be realized in a few years. On the one hand, the simulation suggests that AMS-02 operating for 3 years will measure the positron-to-electron ratio at energy ~ 400 GeV with an error ~ 0.01 [33]. On the other hand, at 400 GeV, the difference of positron-to-electron ratio in the cases with and without primary electron spectrum hardening can be as large as ~ 0.04 (see Fig.2). The detection prospect is thus promising.

---

[1] This conclusion is valid as long as the electron/positron total flux is similar to that measured by Fermi-LAT. If instead, the electron/positron total flux at energies $300-800$ GeV resembles the ATIC data, it is difficult to modify the electron-to-positron ratio unless the primary electron spectrum hardening is much harder than what we have assumed.

We would like to point out that the Australia Telescope National Facility (ATNF) pulsar model also predicts fluctuations of the positron-to-electron ratio and sometimes the ratio drops off quickly [30]. However, unless there is an additional hard electron component, the electron/positron total flux should display the same behavior, which is different from our scenario. The same argument applies to the dark matter model as well.

The spectrum hardening of both primary-electrons and nuclei at ~ 240 GV, if confirmed by AMS-02 in the future, is likely attributed to a "nearby" supernova-remnant-like source with a lifetime (in view of the significant softening of electron/positron total spectrum at the energy $E_{\rm cool} \sim 1$ TeV)

$$\tau \sim 10^{13}~{\rm s}~(\frac{E_{\rm cool}}{1~{\rm TeV}})^{-1}(\frac{u_{\rm tot}}{1~{\rm eV~cm^{-3}}})^{-1}, \qquad (4)$$

where $u_{\rm tot} = u_{\rm B}+u_{\rm cmb}+u_{\rm dust}+u_{\rm star}$, $u_{\rm B} = B_{\rm IG}^2/8\pi$ is the magnetic field energy density, $u_{\rm cmb}$, $u_{\rm dust}$ and $u_{\rm star}$ are the photon energy densities of cosmic microwave background, dust emission and the star emission, respectively. Correspondingly, the source should be within the characteristic cosmic ray diffusion radius

$$\begin{aligned} R_{\rm d} &\sim 2\sqrt{\mathcal{D}\cdot\tau} \\ &\sim 1~{\rm kpc}~(\frac{D_0}{10^{28.7}~{\rm cm^2~s^{-1}}})^{1/2}(\frac{E}{1~{\rm TeV}})^{1/6}(\frac{\tau}{10^{13}~{\rm s}})^{1/2}(5) \end{aligned}$$

where $\mathcal{D} = D_0(E/1~{\rm GeV})^{\delta}$ is the spatial diffusion coefficient which is usually regarded as isotropic [3]. In other words the source should be relatively nearby. There are two candidate supernova remnants, i.e., Geminga and Loop I [29, 34]. Here we do not consider nearby pulsars since they are expected to produce electron/positron pairs rather than mainly electrons. The nearby supernova remnants instead produce "ignorable" amount of positrons. This can be straightforwardly understood as the following. Let us make the"optimistic" estimate, i.e., assuming that the total cosmic ray protons above ~ 240 GeV are dominated by the nearby source. The chance for one high energy proton to produce one positron is $P \sim \sigma_{\rm pp} nc\tau/3 \sim 3.5\times10^{-3}~(n/0.5~{\rm cm^{-3}})(\tau/10^{13}~{\rm s})$, where $\sigma_{\rm pp} \approx 70$ mb is the total cross section of production of pions in proton-proton collision, and $n \sim 0.1-1~{\rm cm^{-3}}$ is the number density of the local interstellar medium. The high energy proton loses about 20% energy in one proton-proton collision, roughly one quarter converts into positron via the decay of the positively charged pion (i.e., $\pi^+ \rightarrow \mu^+ + \nu_\mu \rightarrow e^+ + \nu_{\rm e} + \bar{\nu}_\mu + \nu_\mu$). Hence at the energy of 240 GeV, the positron-to-proton ratio is $\sim P/20^{1.7} \sim 2\times10^{-5}~(n/0.5~{\rm cm^{-3}})(\tau/10^{13}~{\rm s})$, well below the value $\sim 3\times10^{-4}$ inferred from the PAMELA/AMS-02 data [8, 11], where the $E^{-2.7}$-like proton spectrum has been taken into account. If the future AMS-02 data show no evidence for the hardening of the primary electron spectrum at 240 GeV, the cosmic ray sources accounting for the nuclei excesses likely locate at a distance further than $R \sim 1.8~{\rm kpc}~(\frac{D_0}{10^{28.7}~{\rm cm^2~s^{-1}}})^{1/2}(\frac{u_{\rm tot}}{1~{\rm eV~cm^{-3}}})^{-1/2}(\frac{E}{0.24~{\rm TeV}})^{-1/3}$.

IV. CONCLUSION AND DISCUSSION

The accurate positron-to-electron ratio data of AMS-02 confirms the positron excess discovered by PAMELA up to the energy ~ 350 GeV. The dark matter annihilation/decay can account for the positron excess but the Fermi-LAT Galactic diffuse emission data has imposed a tight constraint on such a kind of possibility. The channels of annihilation/decay into $\tau^+\tau^-$ pairs have been excluded. The channel $\chi\chi \rightarrow \mu^+\mu^-$ is possible only for very cored dark matter distribution model (e.g., isothermal-sphere profile) while $\chi \rightarrow \mu^+\mu^-$ is viable (see Fig.1). Alternatively, the positron excess detected by PAMELA/Fermi-LAT/AMS-02 may be dominated by pulsar-like astrophysical sources and the contribution by dark matter may be small but detectable. At energies ≤ 100 GeV, a positron/electron component from either dark matter annihilation/decay or pulsar seems to be favored by the AMS-02 data (see Fig.2). In the case of annihilation of dark matter with a rest mass ~ 100 GeV, the cross section into electron/positron pairs is needed to be $\sim 10^{-26}~{\rm cm^3~s^{-1}}$, consistent with all of the current bounds of the indirect detection measurements.

Whether the primary cosmic ray electron spectrum gets hardened at ~ 240 GeV, like that observed in cosmic-ray-nuclei spectra, is an interesting question. If it does, the subsequent positron-to-electron ratio may get flattened or even decreased, depending on the degree of the primary electron spectrum hardening (see Fig.2). The current AMS-02 data show tentative evidence for flattening at energies above ~ 260 GeV. However the errors are still large and a robust conclusion is not possible. In the future, AMS-02 will precisely measure/separate the cosmic ray electrons and positrons up to ~ 1 TeV, with which the primary electron spectrum hardening hypothesis can be unambiguously tested. The spectrum hardening of both primary-electrons and nuclei at ~ 240 GV, if confirmed by AMS-02 in the future, likely suggests a common physical origin. The electron spectrum softening at ~ 1 TeV would thus point to "nearby" but old supernova-remnant-like source(s) with a lifetime $\lesssim 10^{13}$ s, like Geminga and possibly also Loop I.

Acknowledgments

This work was supported in part by 973 Program of China under grant 2013CB837000, by National Natural Science of China under grant 10925315, and by China Postdoctoral science foundation under grants 2012M521136 and 2012M521137. YZF is also supported by the 100 Talents program of Chinese Academy of Sciences and the Foundation for Distinguished Young

Scholars of Jiangsu Province, China (No. BK2012047).

*Corresponding author.

Electric addresses: fenglei@pmo.ac.cn, bixian85@pmo.ac.cn, hnhe@pmo.ac.cn, tkdong@pmo.ac.cn, yzfan@pmo.ac.cn, chang@pmo.ac.cn

---

[1] V. L. Ginzburg and S. I. Syrovatskii, The Origin of Cosmic Rays. New York: Macmillan (1964).
[2] M. Nagano and A. A. Watson, Rev. Mod. Phys., **72**, 689 (2000).
[3] A. W. Strong, I. V. Moskalenko, and V. S. Ptuskin, Annu. Rev. Nucl. Part. Sci., **57**, 285 (2007).
[4] J. Chang *et al.* [The ATIC Collaboration], Nature, **456**, 362 (2008).
[5] S. Torii *et al.*, arXiv:0809.0760 [astro-ph].
[6] F. Aharonian et al. [H.E.S.S. Collaboration], Astrophys. Astron., **508**, 561 (2009).
[7] A. A. Abdo *et al.* [The Fermi LAT Collaboration], Phys. Rev. Lett., **102**, 181101 (2009).
[8] O. Adriani *et al.*, [The PAMELA Collaboration], Nature, **458**, 607 (2009).
[9] M. Ackermann *et al.* [The Fermi-LAT Collaboration], Phys. Rev. Lett., **108**, 011103 (2012)
[10] Y. Z. Fan, B. Zhang, and J. Chang, Int. J. Mod. Phys. D, **19**, 2011 (2010); T. A. Porter, R. P. Johnson, and P.W. Graham, Annu. Rev. Astron. Astrophys., **49**, 155 (2011); L. Bergström, New. J. Phys. **11**, 105006 (2009); X. G. He, Mod. Phys. Lett. A., **24**, 2139 (2009); M. Boezio, *et al.*, New J. Phys., **11**, 105023 (2009).
[11] M. Aguilar, et al. [The AMS-02 collaboration], Phys. Rev. Lett., **110**, 141102 (2013)
[12] J. P. Wefel, *et al.* [The ATIC Collaboration], International Cosmic Ray Conference, **2**, 31 (2008).
[13] Ahn H S *et al.* [The CREAM Collaboration], Astrophys. J, **714**, L89 (2010).
[14] Adriani O *et al.* [The PAMELA Collaboration], Science, **332**, 69 (2011).
[15] A. E. Vladimirov, G. Jóhannesson, I. V. Moskalenko, and T. A. Porter, Astrophys. J, **752**, 68 (2012); H. X. Cui, L. Feng, X. J. Bi, S. W. Cui, and S. P. Yang, submitted (2013).
[16] V. Ptuskin, V. Zirakashvili, and E. S. Seo, Astrophys. J, **763**, 47 (2013).
[17] V. I. Zatsepin, and N. V. Sokolskaya, Astrophys. Astron., **458**, 1 (2006); Q. Yuan, B. Zhang, and X. J. Bi, Phys. Rev. D, **84**, 043002 (2011).
[18] G. Jungman, M. Kamionkowski and K. Griest, Phys. Rept., **267**, 195 (1996); G. Bertone, D. Hooper, and J. Silk, Phys. Rept., **405**, 279 (2005); D. Hooper and S. Profumo, Phys. Rept., **453**, 29 (2008).
[19] N. Arkani-Hamed, D. P. Finkbeiner, T. Slatyer, and N. Weiner, Phys. Rev. D, **79**, 015014 (2009); X. J. Bi, X. G. He, and Q. Yuan, Phys. Lett. B, **678**, 168 (2009); D. Hooper and K. M. Zurek, Phys. Rev. D, **79**, 103529 (2009); L. Bergstrom, J. Edsjo, and G. Zaharijas, Phys. Rev. Lett., **103**, 031103 (2009).
[20] A. Ibarra, and D. Tran, Phys. Rev. Lett., **100**, 061301 (2008); P. F. Yin et al., Phys. Rev. D, **79**, 023512 (2009); C. R. Chen, F. Takahashi and T. T. Yanagida, Phys. Lett. B, **671**, 71 (2009); E. Nardi, F. Sannino and A. Strumia, J. Cosmol. Astropart. Phys., **0901**, 043 (2009); J. Liu, P. F. Yin and S. H. Zhu, Phys. Rev. D, **79**, 063522 (2009).
[21] M. Cirelli, P. Panci, P. D. Serpico. Nucl. Phys. B, **840**, 284 (2010).
[22] M. Ackermann, *et al.*, Astrophys. J., **761**, 91 (2012).
[23] J. F. Navarro, C. S. Frenk and S. D. M. White, Astrophys. J. **462**, 563 (1996).
[24] K. G. Begeman, A. H. Broeils, and R. H. Sanders, Mon. Not. R. Astron. Soc., **249**, 523 (1991).
[25] A. W. Strong, and I. V. Moskalenko, Astrophys. J. **509** 212 (1998).
[26] M. Ackermann *et al.*, Phys. Rev. D, **82**, 092004 (2010).
[27] L. Feng, Q. Yuan, X. Li, and Y. Z. Fan, Phys. Lett. B, **720**, 1 (2013).
[28] C. S. Shen, Astrophys. J. **162**, L181 (1970); L. Zhang and K. S. Cheng, Astron. Astrophys., **368**, 1063 (2001).
[29] S. Profumo, Central European Journal of Physics, **10**, 1 (2012).
[30] D. Malyshev, I. Cholis, and J. Gelfand, Phys. Rev. D, **80**, 063005 (2009).
[31] D. R. Lorimer, in $YoungNeutronStarsandTheirEnvironments$, edited by F. Camilo & B. M. Gaensler, vol. 218 of IAU Symposium, p. 105. (2004).
[32] J. Liu *et al.*, Phys. Rev. D **85**, 043507 (2012).
[33] J. Casaus, J. Phys.: Conf. Ser. **171**, 012045 (2009).
[34] N. J. Shaviv, E. Nakar, and T. Piran, Phys. Rev. Lett., **103**, 111302 (2009).